

ON MIXING INEQUALITIES: RANK, CLOSURE AND CUTTING PLANE PROOFS

SANJEEB DASH [†] AND OKTAY GÜNLÜK [†]

Abstract. We study the mixing inequalities which were introduced by Günlük and Pochet (2001). We show that a mixing inequality which mixes n MIR inequalities has MIR rank at most n if it is a type I mixing inequality and at most $n - 1$ if it is a type II mixing inequality. We also show that these bounds are tight for $n = 2$.

Given a mixed-integer set $P_I = P \cap Z(I)$ where P is a polyhedron and $Z(I) = \{x \in \mathbb{R}^n : x_i \in \mathbb{Z} \forall i \in I\}$, we define mixing inequalities for P_I . We show that the elementary mixing closure of P with respect to I can be described using a bounded number of mixing inequalities, each of which has a bounded number of terms. This implies that the elementary mixing closure of P is a polyhedron.

Finally, we show that any mixing inequality can be derived via a polynomial length MIR cutting plane proof. Combined with results of Dash (2006) and Pudlák (1997), this implies that there are valid inequalities for a certain mixed-integer set that cannot be obtained via a polynomial-size mixing cutting-plane proof.

Key words. 90C11 Mixed integer programming

1. Introduction. Günlük and Pochet [7] study the polyhedral structure of the so-called *mixing* set

$$S = \left\{ s \in \mathbb{R}_+, z \in \mathbb{Z}^n : s + z_k \geq b_k \quad \text{for } k = 1, \dots, n \right\}$$

where $0 < b_1 < b_2 < \dots < b_n \leq 1$, and show that the following n -term *mixing inequalities* are valid for S :

$$s + b_1 z_1 + \sum_{k=2}^n (b_k - b_{k-1}) z_k \geq b_n, \quad (1.1)$$

$$s + (b_1 + 1 - b_n) z_1 + \sum_{k=2}^n (b_k - b_{k-1}) z_k \geq b_n. \quad (1.2)$$

Note that if $b_n = 1$, inequalities (1.1) and (1.2) are identical. These inequalities are called “mixing” inequalities as they combine, or *mix*, mixed-integer rounding (MIR) inequalities based on individual constraints, namely

$$s + b_k z_k \geq b_k, \quad (1.3)$$

for $k = 1, \dots, n$. Notice that if $n = 1$, the mixing inequality (1.1) becomes the MIR inequality (1.3) and the mixing inequality (1.2) simply becomes the inequality defining S . MIR inequalities were introduced by Nemhauser and Wolsey [8] and they are obtained by applying a simple procedure to any implied (base) inequality. Nemhauser and Wolsey [9] also showed that split cuts, introduced by Cook, Kannan and Schrijver [2], are equivalent to MIR inequalities. The inequalities (1.3) are not the only MIR inequalities for S ; see [5] for a recent study on MIR inequalities.

The original description of the mixing set and the mixing inequalities does not assume $0 < b_i \leq 1$ but for the sake of simplicity, and without loss of generality, we

[†]Mathematical Sciences Department, IBM T. J. Watson Research Center, Yorktown Heights, NY 10598 USA

make this assumption. Notice that in the definition of S , the z_i variables do not have lower bounds and therefore if $b_i \notin (0, 1]$ for some i , it is possible to simply “shift” variable z_i by $\lceil b_i \rceil - 1$ and then replace b_i with $b_i - \lceil b_i \rceil + 1$ to obtain $b_i \in (0, 1]$. The mixing inequalities (1.1) and (1.2) are identical to the ones presented in [7] after this simple transformation. We note that MIR inequalities are known to be invariant under “shifting”, and more generally, under unimodular transformations [5].

Let $I' = \{i_1, \dots, i_t\}$ be a subset of $\{1, \dots, n\}$ and let $proj_{[I']}(S)$ denote the projection of S in the space of the s and z_i variables for $i \in I'$. In other words, $proj_{[I']}(S) = \{s \in \mathbb{R}_+, z \in \mathbb{Z}^{|I'|} : s + z_k \geq b_k, \forall k \in I'\}$. Clearly the mixing inequalities

$$s + b_{i_1} z_{i_1} + \sum_{k=2}^{|I'|} (b_{i_k} - b_{i_{k-1}}) z_{i_k} \geq b_{i_{|I'|}}, \quad (1.4)$$

$$s + (b_{i_1} + 1 - b_{i_{|I'|}}) z_{i_1} + \sum_{k=2}^{|I'|} (b_{i_k} - b_{i_{k-1}}) z_{i_k} \geq b_{i_{|I'|}} \quad (1.5)$$

for $proj_{[I']}(S)$ are also valid for S . We refer to inequality (1.4) as $mix1_{I'}$, and to inequality (1.5) as $mix2_{I'}$, and say that these are $|I'|$ -term mixing inequalities, of type I and type II, respectively. Günlük and Pochet in [7] show that $mix1_{I'}$ and $mix2_{I'}$ are facet-defining for the convex hull of S for all $I' \subseteq \{1, \dots, n\}$, and these inequalities completely describe the convex hull of S .

One of our main contributions in this paper is to show how mixing inequalities can be obtained from the inequalities defining S by repeatedly applying the MIR procedure. We show that the inequality $mix1_{I'}$ has MIR-rank at most $|I'|$ and $mix2_{I'}$ has MIR-rank at most $|I'| - 1$. In addition, for $|I'| = 2$, we show that these bounds can be tight. A more general lower bound on the MIR-rank of these inequalities has recently been given by Dey [6]. We also show that $mix1_{I'}$ and $mix2_{I'}$ have MIR cutting-plane proofs of length $O(|I'|^2)$ from the inequalities defining S .

We define mixing inequalities for general mixed-integer sets in a way that they contain the family of MIR inequalities, and show that some important properties of MIR inequalities extend to these inequalities. In particular, we show that the elementary closure of mixing inequalities is polyhedral. More precisely, we show that the mixing closure of any given mixed integer set can be described using a bounded number of mixing inequalities each of which has a bounded number of terms. This result generalizes the result of Cook, Kannan and Schrijver [2] that the elementary closure of MIR inequalities is polyhedral. Finally, we define cutting-plane proofs for general mixed-integer sets using mixing inequalities, and show, using a recent result by Dash [3], that such cutting-plane proofs have exponential worst-case complexity.

The rest of the paper is organized as follows. In the remainder of this section we first review the MIR inequalities and give some of their well-known properties. We then define mixing inequalities for general mixed-integer sets. We present our results on the rank of mixing inequalities in Section 2. In Section 3 we study mixing

inequalities for general mixed-integer sets and formulate the separation problem as a quadratic mixed-integer program. In Section 4 we define the elementary closure of mixing inequalities for general sets and show that it is polyhedral. Finally, in Section 5, we give a polynomial length MIR cutting-plane proof of mixing inequalities.

1.1. MIR inequalities. Wolsey [11] defines a two variable mixed-integer set $Q = \{s \in \mathbb{R}, z \in \mathbb{Z} : s + z \geq b, s \geq 0\}$ and shows that the *basic mixed-integer inequality*

$$s + \hat{b}z \geq \hat{b} \lceil b \rceil, \quad (1.6)$$

where $\hat{b} = b - (\lceil b \rceil - 1)$ is valid and facet-defining for Q . This observation can be used to generate valid inequalities for a general mixed-integer set $P_I = P \cap Z(I)$ where

$$P = \{x \in \mathbb{R}^n : Ax \geq b\}, \quad Z(I) = \{x \in \mathbb{R}^n : x_i \in \mathbb{Z} \forall i \in I\},$$

and I is a subset of $\{1, \dots, n\}$. We assume that any non-negativity constraints on variables are included in the system $Ax \geq b$.

Let $v = Ax - b$ and note that $v \geq 0$ for all $x \in P$. Assume A has m rows and let $\lambda \in \Gamma$ where

$$\Gamma = \{\lambda \in \mathbb{R}^m : (\lambda A)_i \in \mathbb{Z} \text{ for all } i \in I, (\lambda A)_i = 0 \text{ for all } i \notin I\}.$$

Define λ^+ by $\lambda_i^+ = \max\{\lambda_i, 0\}$. Then the equation $-\lambda v + (\lambda A)x = \lambda b$ is valid for P and so is the inequality

$$(-\lambda)^+ v + (\lambda A)x \geq \lambda b. \quad (1.7)$$

In addition, for all points in P_I , $(-\lambda)^+ v$ is non-negative and $(\lambda A)x$ is integral. Let $\beta = \lambda b - (\lceil \lambda b \rceil - 1)$. The basic mixed-integer inequality implies that $(-\lambda)^+ v + \beta(\lambda A)x \geq \beta \lceil \lambda b \rceil$, or equivalently

$$(-\lambda)^+(Ax - b) + \beta(\lambda A)x \geq \beta \lceil \lambda b \rceil, \quad (1.8)$$

is a valid inequality for P_I . This inequality is the *mixed-integer rounding (MIR)* inequality generated by λ . Inequality (1.7) is called the base inequality of the MIR inequality. See [5] for other ways of defining the MIR inequality, and the equivalence of (1.8) with the definition in [9].

Some well known properties of the MIR inequalities are the following:

1. The MIR inequality (1.8) generated by $-\lambda$ is equivalent to the one generated by λ .
2. If a point $x^* \in P$ violates the MIR inequality (1.8), then
 - (a) x^* satisfies $\lfloor \lambda b \rfloor < (\lambda A)x^* < \lceil \lambda b \rceil$;
 - (b) $\lambda^+ v^* < 1$, where $v^* = Ax^* - b$.

An inequality $cx \geq d$ is called a *split cut* for P with respect to I if $cx \geq d$ is satisfied by points in $P \cap \{\alpha x \leq \gamma\}$ and $P \cap \{\alpha x \geq \gamma + 1\}$, where α, γ are integral and $\alpha_i = 0$ for $i \notin I$. We say that $cx \geq d$ is derived using the *disjunction* $(\alpha x \leq \gamma) \vee (\alpha x \geq \gamma + 1)$. It is known that when $\beta \neq 0$, the inequality (1.8) is a split cut for P derived using the disjunction $(\lambda Ax \leq \lfloor \lambda b \rfloor) \vee (\lambda Ax \geq \lceil \lambda b \rceil)$. In addition, every split cut for P is also an MIR inequality generated by some $\lambda \in \Gamma$ [9].

1.2. Mixing inequalities for general mixed-integer sets. Just as the MIR inequality for P_I can be derived using the set Q and the basic mixed-integer inequality (1.6), we next define mixing inequalities for P_I using S and the mixing inequalities for S . These inequalities contain the MIR inequality as a special case.

Let $K = \{1, \dots, t\}$. Given vectors $\lambda^k \in \Gamma$ for $k \in K$, the MIR inequalities

$$(-\lambda^k)^+(Ax - b) + \beta_k(\lambda^k A)x \geq \beta_k \lceil \lambda^k b \rceil,$$

where $\beta_k = \lambda^k b - (\lceil \lambda^k b \rceil - 1)$, are valid for P_I . Just as inequalities (1.1) and (1.2) are obtained by “mixing” inequalities (1.3), we now derive valid inequalities by mixing the above MIR inequalities. For convenience we collect λ^k as rows of a matrix in increasing order of β_k values. More precisely, we call $\Lambda \in \mathbb{R}^{t \times m}$ a *mixing matrix* for P_I if it satisfies the following conditions:

1. $0 < \beta_1 < \beta_2 < \dots < \beta_t$,
2. $\lambda^k \in \Gamma$ for all $k \in K$.

Given a mixing matrix Λ for P_I , we call the following inequality

$$\sum_{j \in M} (\max_{k \in K} \{-\lambda_j^k\})^+(Ax - b)_j + \sum_{k \in K} (\beta_k - \beta_{k-1})(\lambda^k Ax - \lceil \lambda^k b \rceil + 1) \geq \beta_t \quad (1.9)$$

where $\beta_0 = 0$, a *mixing inequality of type I* generated by Λ . We use $(Ax - b)_j$ to denote the j th row of $(Ax - b)$. Similarly, we define a *mixing inequality of type II* generated by Λ to be

$$\begin{aligned} \sum_{j \in M} (\max_{k \in K} \{-\lambda_j^k\})^+(Ax - b)_j + (\beta_1 + 1 - \beta_t)(\lambda^1 Ax - \lceil \lambda^1 b \rceil + 1) \\ + \sum_{k \in K \setminus \{1\}} (\beta_k - \beta_{k-1})(\lambda^k Ax - \lceil \lambda^k b \rceil + 1) \geq \beta_t \end{aligned} \quad (1.10)$$

To see that mixing inequalities (1.9) and (1.10) are valid for P_I , let $v = Ax - b$ and note that $v \geq 0$ for all $x \in P$. For any matrix $\Lambda \in \mathbb{R}^{t \times m}$, the equation system $(\Lambda A)x - \Lambda v = \Lambda b$ is satisfied by all $x \in P$. Dropping v_j variables with negative coefficients from these equations, one obtains the following valid inequalities

$$\lambda^k Ax + \sum_{j \in M} (-\lambda_j^k)^+ v_j \geq \lambda^k b \quad (1.11)$$

for all $k \in K$. Further relaxing inequality (1.11) we obtain

$$\lambda^k Ax + \sum_{j \in M} \max_{k' \in K} \{(-\lambda_j^{k'})^+\} v_j \geq \lambda^k b \quad (1.12)$$

as valid inequalities for P . Now letting $s = \sum_{j \in M} \max_{k' \in K} \{(-\lambda_j^{k'})^+\} v_j$, and $z_k = \lambda^k Ax - (\lceil \lambda^k b \rceil - 1)$ inequality (1.12) becomes $s + z_k \geq \beta_k$. As Λ is a mixing matrix for P_I , for any given $x \in P_I$, we have the corresponding $s \geq 0$ and $z_k \in \mathbb{Z}$ for all $k \in K$. Furthermore, as $\beta_k \in (0, 1]$ and is strictly increasing, any point $x \in P_I$ can be mapped to a point in $S = \{s \in \mathbb{R}_+, z \in \mathbb{Z}^n : s + z_k \geq \beta_k \forall k \in K\}$. Writing the mixing inequalities (1.1) and (1.2) for S and replacing the surrogate variables with the original ones, one obtains inequalities (1.9) and (1.10).

2. MIR rank of mixing inequalities . The elementary MIR closure of P with respect to I , denoted by $P^{[1]}$, is the set of points in P that satisfy all MIR inequalities that can be generated using the inequalities defining P and the integrality of the variables x_i for $i \in I$. It is known that $P^{[1]}$ is a polyhedral set [2] and therefore it suffices to consider only a finite number of (un-dominated) MIR inequalities to obtain the MIR closure. For any integer $t \geq 2$, let $P^{[t]}$ be the MIR closure of $P^{[t-1]}$ with respect to I . Define $P^{[0]} = P$. We say that a valid inequality for P_I has MIR-rank t , for some integer $t \geq 1$, if it is valid for $P^{[t]}$, but not valid for $P^{[t-1]}$. If a valid inequality is implied by $Ax \geq b$, then it has MIR-rank 0.

2.1. MIR rank of type I mixing inequalities . We next study the MIR rank of type I mixing inequalities and show that the rank of the $|I'|$ -term mixing inequality (1.4) is at most $|I'|$. For simplicity, we use $S^{[t]}$ to denote $S_{LP}^{[t]}$ where S_{LP} stands for the continuous relaxation of S .

THEOREM 2.1. *The $|I'|$ -term mixing inequality $mix1_{I'}$ for S has MIR rank at most $|I'|$.*

Proof. We will prove the following claim: for any $n > 0$, $mix1_{\{1, \dots, n\}}$ is valid for $S^{[n]}$. This would imply that inequality (1.1) for S has MIR rank at most n . and therefore inequality (1.4) for $proj_{[I']}(S)$ has rank at most $|I'|$. As all inequalities defining $proj_{[I']}(S)$ are present in the definition of S , the theorem would follow.

A mixing inequality with only one term (e.g., $mix1_{\{1\}}$) is just an MIR inequality and has MIR rank 1. Assume the claim is true for mixing inequalities with $n-1$ terms. We next show that $mix1_{\{1, \dots, n\}}$ is a split cut for $S^{[n-1]}$ derived from the disjunction:

$$(z_1 \geq z_2) \vee (z_1 \leq z_2 - 1).$$

As every split cut for $S^{[n-1]}$ is also an MIR cut for $S^{[n-1]}$, the claim will follow. More precisely, we will show that $mix1_{\{1, \dots, n\}}$ is a split cut for a set S' defined by the inequalities $s + z_1 \geq b_1$, $s + b_1 z_1 \geq b_1$, $mix1_{\{2, \dots, n\}}$ and $mix1_{\{1, k, \dots, n\}}$ for $k = 3, \dots, n$. Clearly, S' contains $S^{[n-1]}$.

For any point (\bar{s}, \bar{z}) in S' which satisfies the inequality $z_1 \geq z_2$,

$$\bar{s} + b_1 \bar{z}_1 + \sum_{k=2}^n (b_k - b_{k-1}) \bar{z}_k \geq \bar{s} + b_2 \bar{z}_2 + \sum_{k=3}^n (b_k - b_{k-1}) \bar{z}_k \geq b_n.$$

The second inequality above is true as points in S' satisfy $mix1_{\{2, \dots, n\}}$.

We now consider a point (\bar{s}, \bar{z}) in S' which satisfies $z_1 \leq z_2 - 1$.

Case 1: Assume (\bar{s}, \bar{z}) satisfies $\bar{z}_2 \leq \bar{z}_k$ for all $k = 3, \dots, n$. This fact, along with the inequality $z_1 \leq z_2 - 1$, implies that $\bar{z}_1 + 1 \leq \bar{z}_k$ for $k \geq 2$. Therefore

$$\bar{s} + b_1 \bar{z}_1 + \sum_{k=2}^n (b_k - b_{k-1}) \bar{z}_k \geq \bar{s} + b_1 \bar{z}_1 + \sum_{k=2}^n (b_k - b_{k-1}) (\bar{z}_1 + 1) = \bar{s} + b_n \bar{z}_1 + b_n - b_1$$

If $\bar{z}_1 \geq 0$, then using $s + b_1 \bar{z}_1 \geq b_1$, we have

$$\bar{s} + b_n \bar{z}_1 + b_n - b_1 \geq \bar{s} + b_1 \bar{z}_1 + b_n - b_1 \geq b_n$$

and therefore (\bar{s}, \bar{z}) satisfies (1.1).

If, on the other hand, $\bar{z}_1 \leq 0$, then using $s + \bar{z}_1 \geq b_1$, we have

$$\bar{s} + b_n \bar{z}_1 + b_n - b_1 = \bar{s} + \bar{z}_1 - (1 - b_n) \bar{z}_1 + b_n - b_1 \geq \bar{s} + \bar{z}_1 + b_n - b_1 \geq b_n$$

and therefore (\bar{s}, \bar{z}) satisfies (1.1).

Case 2: Assume that (\bar{s}, \bar{z}) satisfies $\bar{z}_2 > \bar{z}_k$ for some $k \in \{3, \dots, m\}$, and let t be the smallest index in $\{3, \dots, m\}$ for which this is true. Then $\bar{z}_k > \bar{z}_t$ for $k = 2, \dots, t-1$. This implies that

$$\bar{s} + b_1 \bar{z}_1 + \sum_{k=2}^n (b_k - b_{k-1}) \bar{z}_k \geq \bar{s} + b_1 \bar{z}_1 + (b_t - b_1) \bar{z}_t + \sum_{k=t+1}^n (b_k - b_{k-1}) \bar{z}_k \geq b_n$$

as the second inequality follows from $mix1_{\{1,t,\dots,n\}}$, which is satisfied by points in S' . \square

We note that there are alternative derivations of $mix1_{\{1,\dots,n\}}$ as a split cut for $S^{[n-1]}$. For example, one can replace inequalities $mix1_{\{1,k,\dots,n\}}$ for $k = 3, \dots, n$ with $mix1_{\{1,\dots,k-1,k+1,\dots,n\}}$ for $k = 2, \dots, n-1$ in the definition of S' above. (To see this, modify the previous proof by assuming in *Case 2* that $\bar{z}_k > \bar{z}_{k+1}$ for some $k \in \{2, \dots, n-1\}$, and by assuming in *Case 1* that $\bar{z}_2 \leq \dots \leq \bar{z}_n$.) This derivation, however, leads to an exponential length MIR cutting plane proof of $mix1_{\{1,\dots,n\}}$, whereas, the first one leads to a polynomial length MIR cutting plane proof (discussed in Section 5).

The previous theorem implies the existence of multipliers λ that can be used to derive $mix1_{\{1,\dots,n\}}$ as an MIR inequality (1.8) from the inequalities defining S' above. In the appendix we explicitly give these multipliers.

We next show that the upper bound on rank can be tight for two-term mixing inequalities of type I. Let

$$T = \left\{ s \in \mathbb{R}_+, z \in \mathbb{Z}^2 : s + z_1 \geq b_1; s + z_2 \geq b_2 \right\}$$

and remember that the 2-term mixing inequality for T is $s + b_1 z_1 + (b_2 - b_1) z_2 \geq b_2$.

THEOREM 2.2. *If $0 < b_1 < b_2 < 1/2$, then the 2-term mixing inequality $mix1_{\{1,2\}}$ for T has MIR rank 2.*

Proof. We will construct a point (s^*, z^*) which satisfies all MIR cuts, but violates $mix1_{\{1,2\}}$. Choose $\delta > 0$ such that $b_2 + 2\delta < 1/2$, and set $z_2^* = 1 - \delta$ and $z_1^* = 1 - 2\delta$.

For any $s^* \geq 0$, $(s^*, z^*) \in T_{LP}$, the LP relaxation of T . We will now choose s^* such that the MIR inequalities $s + b_1 z_1 \geq b_1$ and $s + b_2 z_2 \geq b_2$ are satisfied by (s^*, z^*) , but it violates $mix1_{\{1,2\}}$. Now

$$b_2 - (b_2 - b_1)z_2^* - b_1 z_1^* = b_2(1 - z_2^*) + b_1(z_2^* - z_1^*) = b_2\delta + b_1\delta,$$

and $b_2\delta + b_1\delta$ is greater than

$$b_1(1 - z_1^*) = b_1(2\delta) \text{ and } b_2(1 - z_2^*) = b_2\delta.$$

We choose s^* to be any number less than $b_2\delta + b_1\delta$ and larger than $\max\{b_1(2\delta), b_2\delta\}$. Then (s^*, z^*) is violated by $mix1_{\{1,2\}}$, and satisfies the MIR inequalities above.

Assume that some other MIR inequality is violated by (s^*, z^*) , and assume that this inequality is derived using the multipliers $\lambda = (\lambda_1, \lambda_2)$. Define $v_1^* = s^* + z_1^* - b_1$ and $v_2^* = s^* + z_2^* - b_2$. Then, as $b_2 + 2\delta < 1/2$ implies that $b_2 + 1/2 < 1 - 2\delta = z_1^*$, it follows that $v_1^*, v_2^* > 1/2$. Recall that $\lambda \in \mathbb{Z}^2$, and $\lambda^+ v^* < 1$. As the MIR inequality defined by λ is the same as the MIR inequality defined by $-\lambda$, we can assume that the multiplier with maximum magnitude is positive. The only nonzero λ values satisfying the conditions above and yielding distinct inequalities are $(1, 0)$, $(0, 1)$, and $(-1, 1)$. We constructed (s^*, z^*) so that it satisfied the MIR inequalities obtained with the multiplier vectors $(1, 0)$ and $(0, 1)$. The vector $(-1, 1)$ simply yields inequality $mix2_{\{1,2\}}$ (as proved in Lemma 2.3) which is trivially satisfied by (s^*, z^*) . \square

In a recent paper [6], Dey gives a lower bound on the MIR rank of $mix1_{\{1,\dots,n\}}$ for all n which is a growing function of n . More precisely, he shows that the MIR rank of $mix1_{\{1,\dots,n\}}$ is at least $\lceil \log_2(n+1) \rceil$ when $0 < b_1 < \dots < b_n < 1$. When $n = 2$, his result implies Theorem 2.2.

2.2. MIR rank of type II mixing inequalities . In this section we study the MIR rank of type II mixing inequalities and show that the rank of the $|I'|$ -term mixing inequality (1.5) is at most $|I'| - 1$. We start with studying the set T defined earlier.

LEMMA 2.3. *Inequality $mix2_{\{1,2\}}$ is an MIR inequality for T .*

Proof. We first convert the inequalities defining T to equations by adding non-negative slacks v_1, v_2 as follows:

$$s + z_1 - v_1 = b_1, \quad s + z_2 - v_2 = b_2.$$

Subtracting the first equation from the second, and dropping the term $-v_2$, we get $z_2 - z_1 + v_1 \geq b_2 - b_1$ as a valid inequality for T . Applying (1.6), we obtain $(b_2 - b_1)(z_2 - z_1) + v_1 \geq b_2 - b_1$ as an MIR inequality for T ; substituting out v_1 , we obtain $mix2_{\{1,2\}}$. \square

Note that the derivation above does not use the nonnegativity of s and therefore $mix2_{\{1,2\}}$ is valid for the relaxed mixing set

$$T_R = \left\{ (s, z) \in \mathbb{R} \times \mathbb{Z}^2 : s + z_1 \geq b_1; s + z_2 \geq b_2 \right\}.$$

We will now prove a result on the rank of type II mixing inequalities. The proof will be similar to the proof that $mix1_{\{1,\dots,n\}}$ is an MIR inequality for $S^{[n-1]}$ given after Theorem 2.1.

THEOREM 2.4. *The $|I'|$ -term mixing inequality $mix2_{I'}$ for S has MIR rank at most $|I'| - 1$.*

Proof. As in Theorem 2.1, the current theorem will follow from the following claim: the inequality $mix2_{\{1,\dots,n\}}$ for S is valid for $S^{[n-1]}$ and therefore has MIR rank at most $n - 1$.

We showed that this claim is true for $n = 2$. Assume it is true for $n - 1$; in other words, assume all $n - 1$ term mixing inequalities of type II are valid for $S^{[n-2]}$. Let the inequality $s + z_1 \geq b_1$ be expressed as $s + z_1 - v_1 = b_1$ where $v_1 \geq 0$ is a slack variable, and denote it by inequality (M_1) (as in the Appendix). Also consider the following (type II) mixing inequalities (expressed as equations via slacks)

$$s + (b_1 + 1 - b_n)z_1 + (b_k - b_1)z_k + \sum_{i=k+1}^n (b_i - b_{i-1})z_i - v_k = b_n \quad (M'_k)$$

for $k = 3, \dots, n$ and the type II mixing inequality

$$s + (b_2 + 1 - b_n)z_2 + \sum_{i=3}^n (b_i - b_{i-1})z_i - v_{n+1} = b_n \quad (M'_{n+1})$$

that are valid for $S^{[n-2]}$.

We next define multipliers λ' to obtain a base inequality which yields inequality (1.2) as an MIR inequality from the above inequalities. Let

$$\mu'_k = \begin{cases} (b_2 - b_1)\left(\frac{1}{b_n - b_1}\right) & \text{if } k = 1, \\ (b_2 - b_1)\left(\frac{1}{b_{k-1} - b_1} - \frac{1}{b_k - b_1}\right) & \text{for } k = 3, \dots, n \\ -1 & \text{if } k = n + 1, \end{cases}$$

and note that $\mu'_1 + \sum_{k=3}^n \mu'_k = 1$ and $\sum_{k=3}^p \mu'_k = 1 - (b_2 - b_1)/(b_p - b_1)$ for $p = 3, \dots, n$.

Now consider $\mu'_1(M_1) + \sum_{k=3}^n \mu'_k(M'_k)$ and denote it by

$$\alpha_0 s + \sum_{k=1}^n \alpha_k z_k - \sum_{k=1}^n \mu_k v_k = \beta. \quad (2.1)$$

As discussed in the Appendix, $\beta = b_n - (b_2 - b_1)$, $\alpha_0 = 1$, $\alpha_2 = 0$, and $\alpha_k = b_k - b_{k-1}$ for $k = 3, \dots, n$. In addition,

$$\begin{aligned} \alpha_1 &= \mu'_1 + (b_1 + 1 - b_n) \sum_{k=3}^n \mu'_k = \mu'_1(b_n - b_1) + (b_1 + 1 - b_n) \sum_{k=1}^n \mu'_k \\ &= \frac{b_2 - b_1}{b_n - b_1} (b_n - b_1) + (b_1 + 1 - b_n) = b_2 + 1 - b_n. \end{aligned}$$

Therefore equation 2.1 is the same as

$$s + (b_2 + 1 - b_n)z_1 + \sum_{k=3}^n (b_k - b_{k-1})z_k - \sum_{k=1}^n \mu'_k v_k = b_n - (b_2 - b_1), \quad (2.2)$$

and $\sum_{k=1}^{n+1} \mu'_k (M'_k)$ equals

$$(b_2 + 1 - b_n)(z_1 - z_2) + v_{n+1} - \sum_{k=1}^n \mu'_k v_k = -(b_2 - b_1).$$

Finally, setting $\lambda' = \mu'/(b_2 + 1 - b_n)$, and dropping the variables v_k with negative coefficients in $\sum_{k=1}^{n+1} \lambda'_k (M'_k)$, we get

$$(z_1 - z_2) + v_{n+1}/(b_2 + 1 - b_n) \geq -(b_2 - b_1)/(b_2 + 1 - b_n) \quad (2.3)$$

as a valid inequality for $S^{[n-2]}$. Let γ stand for the right-hand-side of the above inequality. Then $\hat{\gamma} = (b_1 + 1 - b_n)/(b_2 + 1 - b_n)$. Applying the basic mixed-integer inequality we get $(b_1 + 1 - b_n)(z_1 - z_2) + v_{n+1} \geq 0$ as an MIR inequality for $S^{[n-2]}$. Substituting out v_{n+1} from the above expression, we obtain $mix2_{\{1, \dots, n\}}$. \square

3. Separating mixing inequalities. In this section we study the separation problem for mixing inequalities for general mixed-integer sets. We first present bounds on the maximum violation of mixing inequalities and then formulate the associated separation problem as an optimization problem.

3.1. Bounding the violation of mixing inequalities. Consider the mixing set $S = \{s \in \mathbb{R}_+, z \in \mathbb{Z}^t : s + z_k \geq \beta_k \forall k \in K\}$ and let S_{LP} denote its continuous relaxation. For a given $(\bar{s}, \bar{z}) \in S_{LP}$, let the violation of the mixing inequality (1.1) be defined as, $\Delta^1(\bar{s}, \bar{z}) = \beta_t - \bar{s} - \sum_{k=1}^t \delta_k \bar{z}_k$ where $\delta_1 = \beta_1$ and $\delta_k = \beta_k - \beta_{k-1}$ for $k = 2, \dots, t$. Similarly, let the violation of the mixing inequality (1.2) be defined as, $\Delta^2(\bar{s}, \bar{z}) = \beta_t - \bar{s} - \sum_{k=1}^t \epsilon_k \bar{z}_k$ where $\epsilon_1 = \beta_1 + 1 - \beta_t$ and $\epsilon_k = \beta_k - \beta_{k-1}$ for $k = 2, \dots, t$.

LEMMA 3.1. *Let $(\bar{s}, \bar{z}) \in S_{LP}$, then $\Delta^1(\bar{s}, \bar{z}) \leq \frac{1}{2}(1 - \frac{1}{t+1})$ and $\Delta^2(\bar{s}, \bar{z}) \leq \frac{1}{2}(1 - \frac{1}{t})$.*

Proof. As (\bar{s}, \bar{z}) satisfies $\bar{s} + \bar{z}_k - \beta_k \geq 0$ for all $k \in K$, we have

$$\sum_{k=1}^t \delta_k \bar{s} + \sum_{k=1}^t \delta_k \bar{z}_k - \sum_{k=1}^t \delta_k \beta_k \geq 0 \quad (3.1)$$

Define $\delta_{t+1} = 1 - \beta_t$ so that $(1 - \beta_k) = \sum_{j=k+1}^{t+1} \delta_j$ for $k \geq 1$. Adding (3.1) to $\Delta^1(\bar{s}, \bar{z})$, we obtain

$$\Delta^1(\bar{s}, \bar{z}) \leq \beta_t - (1 - \beta_t)\bar{s} - \sum_{k=1}^t \delta_k \beta_k \leq \sum_{k=1}^t \delta_k - \sum_{k=1}^t \delta_k \beta_k = \sum_{k=1}^t \delta_k (1 - \beta_k) = \sum_{k=1}^t \sum_{j=k+1}^{t+1} \delta_k \delta_j.$$

Note that $\sum_{k=1}^{t+1} \delta_k = 1$. We can now rewrite the last term in this expression using the following observation

$$\left(\sum_{k=1}^{t+1} \delta_k\right)^2 = \sum_{k=1}^{t+1} \delta_k^2 + 2 \sum_{k=1}^t \sum_{j=k+1}^{t+1} \delta_k \delta_j \implies \sum_{k=1}^t \sum_{j=k+1}^{t+1} \delta_k \delta_j = \frac{1}{2} \left(1 - \sum_{k=1}^{t+1} \delta_k^2\right). \quad (3.2)$$

For $q \geq 1$, define $w(q) = \min\{\sum_{k=1}^q (\delta_k)^2 : \sum_{k=1}^q \delta_k = 1\}$ and note that $w(q) = 1/q$. Therefore

$$\Delta^1(\bar{s}, \bar{z}) \leq \frac{1}{2} \left(1 - \sum_{k=1}^{t+1} \delta_k^2\right) \leq \frac{1}{2} (1 - w(t+1)) = \frac{1}{2} \left(1 - \frac{1}{t+1}\right).$$

Similarly for $\Delta^2(\bar{s}, \bar{z})$ note that $\bar{s} + \sum_{k=1}^t \epsilon_k \bar{z}_k - \sum_{k=1}^t \epsilon_k \beta_k \geq 0$ as (\bar{s}, \bar{z}) satisfies $\bar{s} + \bar{z}_k - \beta_k \geq 0$ for all $k \in K$ and $\sum_{k=1}^t \epsilon_k = 1$. Adding this expression to $\Delta^2(\bar{s}, \bar{z})$, we obtain

$$\Delta^2(\bar{s}, \bar{z}) \leq \beta_t - \sum_{k=1}^t \epsilon_k \beta_k = \sum_{k=1}^t \epsilon_k (\beta_t - \beta_k) = \sum_{k=1}^{t-1} \epsilon_k (\beta_t - \beta_k) = \sum_{k=1}^{t-1} \sum_{j=k+1}^t \epsilon_k \epsilon_j.$$

Combining the fact that $\sum_{k=1}^t \epsilon_k = 1$ with the observation (3.2) above,

$$\Delta^2(\bar{s}, \bar{z}) \leq \frac{1}{2} \left(1 - \sum_{k=1}^t \epsilon_k^2\right) \leq \frac{1}{2} (1 - w(t)) = \frac{1}{2} \left(1 - \frac{1}{t}\right).$$

□

Using the fact that the validity of mixing inequalities (1.9) and (1.10) for the general mixed integer set P_I was shown by mapping points in P_I to points in the mixing set S , we have the following observation. We define the *violation* of an inequality to be the right-hand-side minus the left-hand-side.

COROLLARY 3.2. *For a given point $\hat{x} \in P$ the violation of any t -term mixing inequality (1.9) is at most $\frac{1}{2} \left(1 - \frac{1}{t+1}\right)$. Similarly, the violation of any t -term mixing inequality (1.10) is at most $\frac{1}{2} \left(1 - \frac{1}{t}\right)$.*

Notice that for $t = 1$ this observation implies that the maximum violation of a type I mixing inequality (1.9) is $1/4$. This is same as the bound shown in [5] for the maximum violation of an MIR inequality. In addition, when $t = 1$, the maximum violation of a type II mixing inequality (1.9) is zero, as the inequality is implied by $Ax \geq b$.

3.2. Separating violated mixing inequalities. For a given a point $\hat{x} \in P$, a most violated mixing inequality (1.9) generated by a mixing matrix that has up to t rows can be obtained by solving the following quadratic mixed-integer program which

we call *Mix-Sep-I*:

$$\text{Maximize} \quad \beta_t - \sum_{j \in M} \delta_j \hat{v}_j - \beta_1 z_1 - \sum_{k \in K \setminus \{1\}} (\beta_k - \beta_{k-1}) z_k$$

Subject to

$$\begin{aligned} \alpha^k &= (\lambda^k A) & k \in K, \\ \alpha_i^k &= 0 & \forall k \in K, i \in N \setminus I, \\ z_k &= (\lambda^k A) \hat{x} - \theta_k & \forall k \in K, \\ \beta_k &= (\lambda^k b) - \theta_k & \forall k \in K, \\ \beta_k &\geq \beta_{k-1} & \forall k \in K, \\ \delta &\geq -\lambda^k & \forall k \in K, \\ 1 &\geq \beta_t, \end{aligned}$$

$$\lambda^k \in \mathbb{R}^m, \alpha^k \in \mathbb{Z}^n \quad k \in K; \theta \in \mathbb{Z}^t, z \in \mathbb{R}^t, \beta \in \mathbb{R}_+^t, \delta \in \mathbb{R}_+^m.$$

where $\hat{v} \in \mathbb{R}^m$ denotes $A\hat{x} - b$. In this formulation, variable z_k stands for $(\lambda^k A \hat{x} - \lceil \lambda^k b \rceil + 1)$ and θ_k stands for $\lceil \lambda^k b \rceil - 1$ (if $\lambda^k b$ is integral, then θ_k can take on the value $\lambda^k b$ or $\lambda^k b - 1$). The objective function measures the violation of the mixing inequality (1.9), defined to be the right-hand-side of the inequality minus the left-hand-side.

LEMMA 3.3. *For a given point $\hat{x} \in P$, an optimal solution of *Mix-Sep-I* corresponds to a most violated mixing inequality of type I that can be generated by a mixing matrix with t or fewer rows.*

Proof. Given a mixing matrix $\Lambda' \in \mathbb{R}^{t' \times m}$ where $t' \leq t$, it is easy to construct a feasible solution to *Mix-Sep-I* where the objective value is the same as the violation of the mixing inequality generated by Λ' . This can simply be done by first appending $t - t'$ copies of the last row of Λ' to obtain the matrix $\Lambda \in \mathbb{R}^{t \times m}$. Letting $\lambda^k = k$ th row of Λ and $\alpha^k = \lambda^k A$ and $\theta_k = \lceil \lambda^k b \rceil - 1$ for $k \in K$, and $\delta_j = (\max_{k \in K} \{-\lambda_j^k\})^+$ for $j \in M$ gives the desired solution to *Mix-Sep-I*.

On the other hand, given an optimal solution to *Mix-Sep-I*, let $\Lambda \in \mathbb{R}^{t \times m}$ be the matrix with k th row equal to the value of λ^k in the solution and note that Λ is not necessarily a mixing matrix as *Mix-Sep-I* does not guarantee that $\beta_k > \beta_{k-1}$ for $k \in K \setminus \{1\}$. Furthermore, the β_k values produced by *Mix-Sep-I* are guaranteed to be equal to $\lambda^k b - (\lceil \lambda^k b \rceil - 1)$ only when $\lambda^k b \notin \mathbb{Z}$. If $\lambda^k b \in \mathbb{Z}$ for some $k \in K$, it is possible that $\theta_k = \lambda^k b$ and $\beta_k = 0$ in the optimal solution. Let Λ' be obtained from Λ by deleting rows λ^k such that $\beta_k = 0$ or $\beta_k = \beta_{k-1}$ in the optimal solution to *Mix-Sep-I*. Notice that Λ' is a mixing matrix with at most t rows. Furthermore, the violation of the mixing inequality (1.9) generated by Λ' equals the optimal value of *Mix-Sep-I*. \square

Similarly, we define *Mix-Sep-II* to be the quadratic mixed-integer program ob-

tained from Mix-Sep-I by changing its objective function to

$$\beta_t - \sum_{j \in M} \delta_j \hat{v}_j - (\beta_1 + 1 - \beta_t)z_1 - \sum_{k \in K \setminus \{1\}} (\beta_k - \beta_{k-1})z_k.$$

LEMMA 3.4. *For a given point $\hat{x} \in P$, an optimal solution of Mix-Sep-II corresponds to a most violated mixing inequality of type II that can be generated by a mixing matrix with t or fewer rows.*

Proof. As in the proof of Lemma 3.3, for a given mixing matrix Λ' with at most t rows, it is easy to construct a feasible solution to Mix-Sep-II with an objective value equal to the violation of the mixing inequality (1.10) generated by Λ' .

Further, for a given optimal solution of Mix-Sep-II let $\Lambda' \in \mathbb{R}^{l \times m}$ be obtained by collecting rows λ^k such that $\beta_k > \beta_{k-1}$ in the optimal solution. Let z^* denote the objective value of this solution. If $\beta_1 > 0$ in the optimal solution, Λ' is a mixing matrix and gives a mixing inequality (1.10) with violation at least z^* . On the other hand, if $\beta_1 = 0$, Λ' is not a mixing matrix, however, it is possible to obtain a new matrix $\bar{\Lambda}$ by moving the first row of Λ' to the end. The matrix $\bar{\Lambda}$ gives $\bar{\lambda}^1 b - (\lceil \bar{\lambda}^1 b \rceil - 1) > 0$ and $\bar{\lambda}^l b - (\lceil \bar{\lambda}^l b \rceil - 1) = 1$. If $\bar{\lambda}^{l-1} b - (\lceil \bar{\lambda}^{l-1} b \rceil - 1)$ is also 1, then we further delete the last row of $\bar{\Lambda}$ to obtain a mixing matrix where the violation of the associated mixing inequality (1.10) equals z^* . \square

4. Mixing closure of mixed-integer sets. We define the *mixing closure* of P with respect to I to be the set of points in P that satisfy all mixing inequalities (1.9) and (1.10) that can be generated by mixing matrices. Let $clo(P_I)$ denote the mixing closure of P with respect to I . Our main result in this section is that $clo(P_I)$ can be described using a bounded number of mixing inequalities each of which has a bounded number of terms. In other words, it is sufficient to consider a bounded number of mixing matrices, each having a bounded number of rows. As we only consider rational data, without loss of generality, we assume that (after scaling, if necessary) $A \in \mathbb{Z}^{m \times n}$ and $b \in \mathbb{Z}^m$ in the definition of P . Before presenting our main result, we first study a special case where all variables are integral.

4.1. Mixing closure of pure integer sets. It is significantly easier to analyze $clo(P_I)$ when there are no continuous variables in the definition of P_I ; that is, when $I = N$. Let $\Lambda \in \mathbb{R}^{t \times m}$ be a mixing matrix for P_N and consider the mixing inequality of type I generated by Λ

$$\sum_{j \in M} \delta_j v_j + \sum_{k \in K} (\beta_k - \beta_{k-1})(\lambda^k A x - \lceil \lambda^k b \rceil + 1) \geq \beta_t \quad (4.1)$$

where $v = Ax - b$ and $\delta_j = (\max_{k \in K} \{-\lambda_j^k\})^+$. We next observe that it is sufficient to consider mixing matrices with small entries.

LEMMA 4.1. *Let $\bar{x} \in P$ and $\Lambda \in \mathbb{R}^{t \times m}$ be a mixing matrix for P_N . If \bar{x} violates a type I mixing inequality (4.1) generated by Λ then there exists a mixing matrix*

$\Lambda' \in \mathbb{R}^{t \times m}$ for P_N such that $\mathbf{1} > \Lambda' > -\mathbf{1}$ and \bar{x} also violates the mixing inequality generated by Λ' .

Proof. Assume that $\delta_l > 0$ for some $l \in M$. In other words $0 > \min_{k \in K} \{\lambda_l^k\}$. Consider Λ' obtained by replacing λ_l^k with $\lambda_l^k + \lfloor \delta_l \rfloor$ for all $k \in K$. Clearly Λ' is a mixing matrix. The left-hand-side of the mixing inequality generated by Λ' is

$$\sum_{j \in M} \delta_j \bar{v}_j + \sum_{k \in K} (\beta_k - \beta_{k-1}) (\lambda^k A \bar{x} - \lceil \lambda^k b \rceil + 1) - \lfloor \delta_l \rfloor \bar{v}_l + \sum_{k \in K} (\beta_k - \beta_{k-1}) \lfloor \delta_l \rfloor (a_l \bar{x} - b_l)$$

where $\bar{v}_j = a_j \bar{x} - b_j \geq 0$ is the surplus variable associated with the j th row. Note that the right-hand-side of the inequality is the same as the right-hand-side of inequality (4.1) as A and b are integral. Using $1 \geq \beta_t = \sum_{k=1}^t (\beta_k - \beta_{k-1})$ and $\bar{v}_l, \lfloor \delta_l \rfloor \geq 0$ we have

$$- \lfloor \delta_l \rfloor \bar{v}_l + \sum_{k \in K} (\beta_k - \beta_{k-1}) \lfloor \delta_l \rfloor (a_l \bar{x} - b_l) = - \lfloor \delta_l \rfloor \bar{v}_l + \beta_t \lfloor \delta_l \rfloor \bar{v}_l \leq - \lfloor \delta_l \rfloor \bar{v}_l + \lfloor \delta_l \rfloor \bar{v}_l = 0.$$

Therefore, the mixing inequality generated by Λ' is violated at least as much as the original inequality (4.1). Without loss of generality, we can therefore assume that $\Lambda' > -\mathbf{1}$.

Now assume $\lambda_l^i \geq 1$ for some $l \in M$ and some $i \in K$ and consider Λ' obtained by replacing λ_l^i with $\lambda_l^i - \lfloor \lambda_l^i \rfloor$. Λ' is a mixing matrix and the left-hand-side of the mixing inequality generated by Λ' is

$$\sum_{j \in M} \delta_j \bar{v}_j + \sum_{k \in K} (\beta_k - \beta_{k-1}) (\lambda^k A \bar{x} - \lceil \lambda^k b \rceil + 1) - (\beta_i - \beta_{i-1}) \lfloor \lambda_l^i \rfloor (a_l \bar{x} - b_l)$$

and the right-hand-side is the same as inequality (4.1) as the data is integral. Clearly the new inequality is violated at least as much as the original inequality (4.1) as $(\beta_i - \beta_{i-1}) \lfloor \lambda_l^i \rfloor \geq 0$ and $a_l \bar{x} \geq b_l$. Therefore all $\lambda_l^i \geq 1$ can be replaced with $\lambda_l^i - \lfloor \lambda_l^i \rfloor$ to obtain a mixing matrix $\Lambda' < \mathbf{1}$. \square

Based on this observation, we next show that there are a finite number of mixing matrices for P_N and therefore the elementary closure of mixing inequalities of type I is polyhedral. Let $\Delta \in \mathbb{Z}_+$ denote the absolute value of the largest entry in $[A, b]$ and let $t^* = (2m\Delta)^{(n+1)}$.

LEMMA 4.2. *If $\bar{x} \in P$ violates a type I mixing inequality (4.1) then it violates one with at most t^* terms.*

Proof. By definition $[A, b] \in [-\Delta, \Delta]^{m \times (n+1)}$. Using Lemma 4.1, and without loss of generality, we can therefore assume that if $\bar{x} \in P$ violates a mixing inequality with t terms, then it violates one generated by a mixing matrix that satisfies $(\Lambda A, \lfloor \Lambda b \rfloor) \in (-m\Delta, m\Delta)^{t \times (n+1)}$. Therefore, in Mix-Sep-I it suffices to consider only $\kappa = (2m\Delta)^{t \times (n+1)}$ possible choices for variables (α, θ) .

In addition, note that for any $\bar{x} \in P$, it suffices to consider mixing inequalities with at most t^* terms as the term $q_k \stackrel{\text{def}}{=} \lambda^k A \bar{x} - \lceil \lambda^k b \rceil + 1$ in inequality (4.1) can be assumed to be strictly increasing and there are only t^* possible choices for $(\lambda^k A, \lceil \lambda^k b \rceil - 1)$.

Given any violated mixing inequality, if $q_k \geq q_{k+1}$ then one can throw away the term q_k for $k > 1$ and replace the coefficient of q_{k+1} with $\beta_{k+1} - \beta_{k-1}$ to obtain a mixing inequality with fewer terms and at least as much violation. \square

Let $clo^1(P_I)$ denote the set of points in P that satisfy all mixing inequalities of type I that can be generated by mixing matrices. Define $clo^2(P_I)$ similarly using mixing inequalities of type II. We next observe that $clo^1(P_I)$ is polyhedral.

COROLLARY 4.3. *$clo^1(P_N)$ is a polyhedron.*

Proof. Using Lemma 4.2, it suffices to consider at most $\kappa^* = (2m\Delta)^{t^* \times (n+1)}$ possible choices for (α, θ) in Mix-Sep-I to obtain a violated mixing inequality of type I. Notice that after fixing (α, θ) , the value of the z variables are implied and therefore, for each fixed value of (α, θ) , the most violated inequality can be obtained by solving a linear program obtained from Mix-Sep-I by fixing α, θ and z variables. As it is sufficient to consider the basic feasible solutions when solving a linear program, and as there a finite number of such basic feasible solutions, say w^* , one only needs to consider $w^* \kappa^*$ inequalities to obtain a violated one. \square

Note that for each mixing matrix Λ it is possible to write a type I mixing inequality and a type II mixing inequality. In other words, Mix-Sep-I and Mix-Sep-II have identical feasible regions and only differ in their objective functions. Using this basic observation, it is possible to adopt Lemmas 4.1 and 4.2 to mixing inequalities of type II and show that $clo^2(P_N)$ and therefore $clo(P_N) = clo^1(P_N) \cap clo^2(P_N)$ is a polyhedron. As our results in the next section subsume this result, we do not present it and avoid repetition.

4.2. Mixing closure of mixed-integer sets. In this section we show that $clo(P_I)$ is a polyhedron. Unlike the pure integer case ($I = N$), we are not able to show that $clo(P_I)$ is given by mixing matrices with small entries. We instead argue that it suffices to consider mixing matrices with “bounded fractionality”, i.e., matrices whose entries are integer multiples of some rational number that depends on A and b . We also argue that fractionality of the coefficients β_i ($i = 1, \dots, t$) in a non-redundant mixing inequality is also bounded and therefore it suffices to consider mixing inequalities with a bounded number of terms. Using these observations, we then show that $clo(P_I)$ is a polyhedron. This result is motivated by a similar result for non-redundant MIR cuts in [5], but the proof is substantially more complicated.

Remember that, without loss of generality, A and b are assumed to be integral. Let $g(A)$ stand for the maximum subdeterminant of A , and let $f(A)$ stand for the product of distinct subdeterminants of A . Clearly, $f(A)$ is a divisor of $g(A)!$.

One can obtain trivial upper bounds for $g(A)$ and $f(A)$ as follows. For a square $t \times t$ matrix B with columns b_1, \dots, b_t , $det(B) \leq \prod_{i=1}^t \|b_i\| \leq (\sqrt{t} \max_{i,j} |B_{ij}|)^t$. For positive integers k, q , define $h(k, q) = (\sqrt{kq})^k$. Then $g(A) \leq h(\min\{m, n\}, \max_{i,j} |A_{ij}|)$, and

$$f(A) \text{ is a divisor of } h(\min\{m, n\}, \max_{i,j} |A_{ij}|)!$$

defined by p linearly independent constraints, and corresponds to a $p \times p$ submatrix of \mathcal{A} , say \mathcal{B} (which we refer to as a *basis matrix*). For each column in \mathcal{B} , the corresponding constraint is satisfied as an equation by the basic feasible solution, and we will say that constraint is *present* in \mathcal{B} .

We will prove that a basic optimal solution of Mix-Sep-LP-I defines a mixing inequality with the properties stated in the theorem. To prove this, we will show that all components of the inverse of an optimal basis are integral multiples of $1/\Omega f(A)^2$. This will imply that in a basic optimal solution, the components of β and Λ are integral multiples of $1/\Omega f(A)^2$, as the right-hand-side of Mix-Sep-LP-I is integral.

Consider a basic optimal solution $\mathcal{X}' = (\Lambda', \beta', \delta')$ of Mix-Sep-LP-I with associated basis matrix \mathcal{B} . It defines a mixing inequality, say \mathcal{I}' , with violation at least Δ . Further, it satisfies $0 < \beta'_1 < \beta'_2 < \dots < \beta'_t \leq 1$, otherwise there exists a mixing inequality having fewer than t terms and violation $\geq \Delta$, a contradiction to the minimality of \mathcal{I} . Therefore, out of the last $3t - 1$ columns of \mathcal{A} , only the last one (corresponding to $\beta_t \leq 1$) can be present in \mathcal{B} . If any of the other $3t - 2$ columns is present in \mathcal{B} , then one of the following constraints is satisfied by β' as an equation: $\beta_{i+1} - \beta_i \geq 0$ for $i = 1, \dots, t - 1$, or $\beta_i \geq 0$ for $i = 1, \dots, t$ or $\beta_i \leq 1$ (or $-\beta_i \geq -1$) for $i = 1 \dots t - 1$.

This implies that the columns corresponding to the constraints $\lambda^i b - \beta_i = \lceil \lambda^i b \rceil - 1$ for $i = 1, \dots, t - 1$ must be present in \mathcal{B} : if any column (say the i th one) is absent, then the $(tm + m + i)$ th row of \mathcal{B} does not have any nonzero entries and \mathcal{B} does not have full row rank, a contradiction. Finally, at least one of the constraints $\beta_t \leq 1$ and $\lambda^t b - \beta_t = \lceil \lambda^t b \rceil - 1$ is present in \mathcal{B} . We now assume that the constraints involving β present in \mathcal{B} are permuted to the end of \mathcal{B} .

Case 1: If only one of the constraints $\beta_t \leq 1$ and $\lambda^t b - \beta_t = \lceil \lambda^t b \rceil - 1$ is present in \mathcal{B} , it has the form

$$\mathcal{B} = \begin{bmatrix} M & B \\ 0_5 & -I_t \end{bmatrix} \Rightarrow \mathcal{B}^{-1} = \begin{bmatrix} M^{-1} & M^{-1}B \\ 0_5 & -I_t \end{bmatrix}.$$

Here 0_5 is a $t \times (tm + m)$ matrix with zero entries, M is a nonsingular square matrix with $tm + m$ rows and B is a matrix with nonzero components drawn from the vector b (and thus has only integral entries).

Case 2: If both the constraints $\beta_t \leq 1$ and $\lambda^t b - \beta_t = \lceil \lambda^t b \rceil - 1$ are present in \mathcal{B} , it has the form

$$\mathcal{B} = \begin{bmatrix} M' & 0_6 & B' \\ 0_7 & -e_t & -I_t \end{bmatrix},$$

where M' has $tm + m$ rows but $tm + m - 1$ columns, 0_6 and 0_7 are matrices of appropriate dimension with zero entries, the $(tm + m)$ th column corresponds to $\beta_t \leq 1$, and subsequent columns correspond to $\lambda^i b - \beta_i = \lceil \lambda^i b \rceil - 1$ for $i = 1, \dots, t$. Let \mathcal{B}' be obtained from \mathcal{B} by subtracting the p th column from the $(tm + m)$ th column. Then $\mathcal{B}' = \mathcal{B} \times T$, where T is a $p \times p$ matrix with $T_{ij} = 1$ if $i = j$, or $T_{ij} = -1$

if $(i, j) = (p, tm + m)$, and 0 otherwise. Then $\mathcal{B}^{-1} = T(\mathcal{B}')^{-1}$. Notice that \mathcal{B}' has the same block upper triangular structure as \mathcal{B} in Case 1. We will therefore focus on analyzing the components of the inverse (especially their denominators) of a basis matrix having the form in Case 1.

Let the columns of M corresponding to the constraints involving λ^i be M_i for $i = 1, \dots, t$. Let N_i be the submatrix of M_i obtained by choosing the m rows corresponding to the variables λ^i ; clearly N_i is a submatrix of $[A I_m]$ for $i = 1, \dots, t - 1$, and a submatrix of $[A b I_m]$ for $i = t$ (the vector b is present in N_t only in the matrix \mathcal{B}' in Case 2). Further, the only nonzeros in the m rows of M corresponding to λ^i are contained in N_i ; as these rows are linearly independent, N_i has rank m , and at least m columns. Also, the columns of $[A b]$ present in N_i are linearly independent, as the columns in M_i are linearly independent, for $i = 1, \dots, t$ (see the depiction of M_i below). We can combine the above facts to conclude that N_i contains a nonsingular $m \times m$ submatrix A_i containing all columns from $[A b]$ present in N_i . Then the columns in N_i but not in A_i (denote these by $N_i \setminus A_i$) are unit vectors, and correspond to constraints $\lambda_j^i + \delta_j \geq 0$ for different j . We depict M_i below, and its various submatrices which we refer to in this proof. Assume M_i has $m + l$ columns for some $l \geq 0$, and A_i has k columns of A for some $k \leq m$. Let a_i be the i th column of A , for $i = 1, \dots, n$, and let i_1, \dots, i_k be distinct integers in $[1, n]$, and let i_{k+1}, \dots, i_{m+l} be distinct integers in $[1, m]$. Then M_i has the form below:

$$M_i \rightarrow \left[\begin{array}{c} 0_{m(i-1) \times (m+l)} \\ \left[\begin{array}{cc} A_i & N_i \setminus A_i \\ \boxed{a_{i_1} \dots a_{i_k} e_{i_{k+1}} \dots e_{i_m}} & \boxed{e_{i_{m+1}} \dots e_{i_{m+l}}} \end{array} \right] N_i \\ 0_{m(t-i) \times (m+l)} \\ \underbrace{\boxed{0_{m \times k} e_{i_{k+1}} \dots e_{i_m}}}_{B_i} e_{i_{m+1}} \dots e_{i_{m+l}} \end{array} \right];$$

where e_j stands for a unit vector in \mathbb{R}^m with a one in the j th position and zeros elsewhere.

Let $M(A_i)$ stand for the columns of M_i which intersect A_i and $M(N_i \setminus A_i)$ stand for the remaining columns in M_i . Let the columns $M(N_i \setminus A_i)$ be arranged at the end of M , for $i = 1, \dots, t$. Then M is a non-singular block arrow matrix having the following form:

$$M = \begin{bmatrix} A_1 & \dots & 0 & C_1 \\ & & \dots & \\ 0 & \dots & A_t & C_t \\ B_1 & \dots & B_t & D \end{bmatrix}.$$

Here M has $(t + 1)^2$ blocks of $m \times m$ matrixes, where the diagonal blocks are A_1, \dots, A_t, D (we will describe D in a moment). The blocks in the last row (other

than D) are B_1, \dots, B_t . Each block B_i is a square submatrix of $M(A_i)$ with m rows corresponding to the variables $\delta_i (i = 1, \dots, m)$; some of its columns are distinct unit vectors, and the remaining columns have only zero entries (see the depiction of M_i above). Each block C_i is an $m \times m$ matrix and consists of the columns of $N_i \setminus A_i$ along with columns with zero entries. As discussed above, the nonzero columns of C_i are distinct unit vectors. Further, the columns of C_i and C_j for $i \neq j$ have nonzeros in non-overlapping columns. We can conclude that at most m of the blocks C_i are nonzero. Finally, each column of D is a unit vector; either it corresponds to a constraint $\delta_j \geq 0$ or $\lambda_j^i + \delta_j \geq 0$ for some $j \in [1, m]$.

The inverse of M is not hard to compute. We start off with the LU decomposition of M :

$$LU = M \implies L = \begin{bmatrix} I_m & \dots & 0 & 0 \\ 0 & \dots & I_m & 0 \\ B_1 A_1^{-1} & \dots & B_t A_t^{-1} & I_m \end{bmatrix}, U = \begin{bmatrix} A_1 & \dots & 0 & C_1 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & A_t & C_t \\ 0 & \dots & 0 & \bar{D} \end{bmatrix},$$

where $\bar{D} = D - \sum_{i=1}^t B_i A_i^{-1} C_i$. The products $B_i A_i^{-1} C_i \neq 0$ for only m distinct values of i ; without loss of generality, we assume that $\bar{D} = D - \sum_{i=1}^m B_i A_i^{-1} C_i$. As M is non-singular, so is \bar{D} . Further, as the unit vectors in C_i and C_j for $i \neq j$ are in non-overlapping columns, and as the nonzero rows of B_i are unit vectors, the nonzero entries in $\bar{D} - D$ are simply components of A_i^{-1} , and are thus ratios of subdeterminants of $[A b]$. This implies that every entry of $\bar{D} - D$ is an integral multiple of $1/f(A)$. As the components of D are either zero or one, every component of $f(A)\bar{D}$ is integral and is bounded in magnitude by $f(A) + f(A)g(A) = f(A)(1 + g(A))$ which is a divisor of $(g(A) + 1)!$. Therefore every component of $(f(A)\bar{D})^{-1}$ is an integral multiple of $1/f(A)\bar{D}$ which is an integral multiple of $1/\Omega = 1/h(m, (g(A) + 1)!)!$. Therefore every component of $\bar{D}^{-1} = f(A)^m (f(A)\bar{D})^{-1}$ is an integral multiple of $1/\Omega$.

Finally,

$$L^{-1} = \begin{bmatrix} I_m & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & I_m & 0 \\ -B_1 A_1^{-1} & \dots & -B_t A_t^{-1} & I_m \end{bmatrix}, U^{-1} = \begin{bmatrix} A_1^{-1} & \dots & 0 & -A_1^{-1} C_1 \bar{D}^{-1} \\ \dots & \dots & \dots & \dots \\ 0 & \dots & A_t^{-1} & -A_t^{-1} C_t \bar{D}^{-1} \\ 0 & \dots & 0 & \bar{D}^{-1} \end{bmatrix}$$

Clearly every component of L^{-1} is an integral multiple of $1/f(A)$. Every component of U^{-1} is an integral multiple of $1/\Omega f(A)$. Therefore every component of $M^{-1} = U^{-1} L^{-1}$ is an integral multiple of $1/\Omega f(A)^2$. As the matrix B' in \mathcal{B} has integral entries, and because of the relationship between \mathcal{B}^{-1} and M^{-1} , every entry in \mathcal{B}^{-1} is an integral multiple of $1/\Omega f(A)^2$ (this is also true in Case 2 above).

As the right-hand-side of Mix-Sep-LP-I has integral components, it follows that the components of $\mathcal{X}' = (\Lambda', \beta', \delta')$ are all integral multiples of $1/\Omega f(A)^2$. Further, as the associated mixing inequality \mathcal{I}' has distinct values of β_i s contained in the interval $(0, 1]$, it follows that \mathcal{I}' has at most $\Omega f(A)^2$ terms. \square

COROLLARY 4.5. *Let $\bar{x} \in P$ and assume it violates a type II mixing inequality (1.10). Then \bar{x} violates a type II mixing inequality with at most $\Omega f(A)^2$ terms such that each β_i is an integral multiple of $1/\Omega f(A)^2$.*

Proof. As in the proof of Theorem 4.4, consider the collection of violated type II mixing inequalities for \bar{x} and from among them let \mathcal{J} be one that has fewest number of terms. Let the violation of \mathcal{J} be $\Delta > 0$ and assume that it is generated by the mixing matrix $\bar{\Lambda} \in \mathbb{R}^{t \times m}$. Let $\bar{z} = \bar{\Lambda}A\bar{x} - \lceil \bar{\Lambda}b \rceil + 1$ and $\bar{v} = A\bar{x} - b$. Recall that Mix-Sep-LP-II is the linear program defined by optimizing

$$\max \beta_t - \sum_{i=1}^m \delta_i \bar{v}_i - (\beta_1 + 1 - \beta_t) \bar{z}_1 - \sum_{i=2}^t (\beta_i - \beta_{i-1}) \bar{z}_i$$

subject to the constraints of Mix-Sep-LP-I, i.e., to (4.3) - (4.7). As \mathcal{J} corresponds to a solution of Mix-Sep-LP-II, an optimal solution of Mix-Sep-LP-II defines a violated mixing inequality of type II with violation at least Δ .

Consider a basic optimal solution of Mix-Sep-LP-II, with associated mixing inequality \mathcal{J}' . If it satisfies $\beta_i = \beta_{i+1}$ for $1 \leq i \leq t-1$, then there exists another violated mixing inequality of type II with fewer terms than \mathcal{J} , a contradiction. In addition, if $\beta_0 = 0$ and $\beta_t = 1$, then, as discussed in the proof of Lemma 3.4, there exists a mixing matrix, and a corresponding type II mixing inequality with violation Δ and $\beta_0 > 0$ and $\beta_t = \beta_{t-1} = 1$, again a contradiction.

Therefore, we can assume that any basic optimal solution satisfies $\beta_1 < \beta_2 < \dots < \beta_t$ and at most one of $\beta_0 = 0$ and $\beta_t = 1$ holds. In the proof of Theorem 4.4 we showed that any such basic feasible solution of Mix-Sep-LP-I has the property that β_1, \dots, β_t are integral multiples of $1/\Omega f(A)^2$. As the basic solution which yields \mathcal{J}' is a basic feasible solution of Mix-Sep-LP-I satisfying the above condition on the β_i s, we can conclude that the β_i values in \mathcal{J}' are integral multiples of $1/\Omega f(A)^2$. \square

To prove that $\text{clo}(P_I)$ is a polyhedron, we will use a proof technique similar to the one in [4] used for showing that the MIR closure of P with respect to I is a polyhedron.

THEOREM 4.6. *The mixing closure of P with respect to I is a polyhedron.*

Proof. Let $q = \Omega f(A)^2$. Define

$$C = \{\beta \in \mathbb{R}^q : 0 \leq \beta_1 \leq \dots \leq \beta_q \leq 1, \beta_i \text{ is an integral multiple of } 1/q, \text{ for } i = 1, \dots, q\}.$$

C is clearly a finite set. For some vector $\bar{\beta} \in C$, define Mix-Sep-I($\bar{\beta}$) to be the integer program obtained by fixing the values of β_i in Mix-Sep-I to $\bar{\beta}_i$; notice that the objective function of Mix-Sep-I($\bar{\beta}$) is a linear function of the variables. The convex hull of solutions of this integer program (call it the *integer hull*) has finitely many vertices. Define Mix-Sep-II($\bar{\beta}$) in a similar manner.

Given a point $\bar{x} \in P \setminus \text{clo}(P_I)$, Theorem 4.4 implies that there exists a violated mixing inequality which defines a solution of Mix-Sep-I($\bar{\beta}$) or Mix-Sep-II($\bar{\beta}$) for some $\bar{\beta} \in C$. Therefore, there exists a violated mixing inequality associated with a vertex

of the integer hull of $\text{Mix-Sep-I}(\bar{\beta})$; note that $\text{Mix-Sep-II}(\bar{\beta})$ has the same integer hull. This implies that $\text{clo}(P_I)$ is the set of points satisfying the mixing inequalities associated with the vertices of the integer hull of $\text{Mix-Sep-I}(\bar{\beta})$ for all $\bar{\beta} \in C$. Therefore $\text{clo}(P_I)$ is a polyhedron. \square

5. Lengths of MIR proofs for mixing inequalities . Let $cx \geq d$ be a valid inequality for P_I . An *MIR cutting-plane proof* (or MIR proof) of $cx \geq d$ from P with respect to I is a sequence of inequalities $a_i x \geq d_i$ ($i = 1, \dots, L$) such that the last inequality in the sequence is $cx \geq d$, and for $i = 1, \dots, L$, the inequality $a_i x \geq d_i$ is an MIR inequality derived from the previous inequalities in the sequence and the inequalities in $Ax \geq b$. The *length* of this proof is said to be L . Cutting plane proofs for Gomory-Chvátal cuts or lift-and-project cuts are defined similarly where each inequality in the sequence is required to be a Gomory-Chvátal or lift-and-project cut, respectively, obtained using the previous inequalities in the sequence and $Ax \geq b$, see [3, 10]. Cutting-plane proofs were introduced by Chvátal in [1].

Pudlák in [10] showed that there are valid inequalities for a particular mixed-integer set P_I (arising from a graph problem) that cannot have a polynomial-length Gomory-Chvátal cutting-plane proof. Later Dash [3] showed that the same inequalities cannot have a polynomial-length MIR cutting-plane proof either. In other words, for these particular inequalities, any MIR cutting-plane proof has exponential length.

In this section, we show that the same negative result holds for mixing inequalities. We define a *mixing cutting-plane proof* the same way as above where each inequality in the cutting plane proof is now derived from previous inequalities via mixing as in (1.9) and (1.10). We first show that mixing inequalities (1.1) and (1.2) have an MIR proof of length $O(n^2)$ from S_{LP} . An immediate consequence of this result is that mixing inequalities (1.9) and (1.10) for P_I with t terms have $O(t^2)$ length MIR proofs from $Ax \geq b$. These observations, when combined with results in [3], imply that mixing proofs have exponential encoding size for Pudlák's inequality system.

THEOREM 5.1. *The inequalities $\text{mix}1_{\{1, \dots, n\}}$ and $\text{mix}2_{\{1, \dots, n\}}$ have MIR proofs of length $O(n^2)$ from the set S .*

Proof. For $1 \leq i < j \leq n$, let $\text{ineq}(i, j)$ denote the mixing inequality $\text{mix}1_{\{i, j, j+1, \dots, n\}}$. In Section 2.1 we showed that $\text{mix}1_{\{1, \dots, n\}}$ can be derived as an MIR inequality using inequalities $s + z_1 \geq b_1$, $s + b_1 z_1 \geq b_1$, $\text{ineq}(2, 3)$ and $\text{ineq}(1, k)$ for $k = 3, \dots, n$. It is easy to see that this also implies that any mixing inequality $\text{ineq}(i, j)$ can be derived as an MIR inequality using inequalities $s + z_i \geq b_i$, $s + b_i z_i \geq b_i$ together with $\text{ineq}(j, j+1)$ and $\text{ineq}(i, k)$ for $k = j+1, \dots, n$.

Note that for any $1 \leq i < j \leq n$, inequality $\text{ineq}(i, j)$ has $n - j + 3$ terms and it is derived using mixing inequalities with fewer terms. Based on this observation, it is possible to produce a short MIR cutting-plane proof as follows: First generate all simple MIR inequalities $s + b_i z_i \geq b_i$ for $i = 1, \dots, n$. Next generate all mixing inequalities $\text{ineq}(i, j)$ with 3 terms using base inequalities $s + z_i \geq b_i$ and simple MIR inequalities $s + b_i z_i \geq b_i$. Finally, for all $k = 4, \dots, n$ generate all k -term mixing

inequalities $ineq(i, j)$ using the base inequities, simple MIR inequalities and mixing inequalities $ineq(i, j)$ with $k - 1$ or fewer terms. Notice that all mixing inequalities $ineq(i, j)$ with $k - 1$ or fewer terms are generated before any mixing inequality $ineq(i, j)$ with k or more terms.

Clearly, this procedure produces n simple MIR inequalities and $(n^2 - n)/2$ mixing inequalities and therefore the MIR proof of $mix1_{\{1, \dots, n\}}$ has length at most $O(n^2)$.

An MIR proof of $mix2_{\{1, \dots, n\}}$ with length $O(n^2)$ is derived in a similar manner by defining $ineq'(i, j)$ to denote the mixing inequality $mix2_{\{i, j, j+1, \dots, n\}}$. \square

COROLLARY 5.2. *Mixing proofs have exponential worst-case encoding size.*

Proof. Given a mixing proof of $cx \geq d$ from $Ax \geq b$ of length L , assume the i th mixing inequality in the proof has t_i terms. It follows from Theorem 5.1 that there is an MIR proof of $cx \geq d$ from $Ax \geq b$ with length $O(\sum_{i=1}^L t_i^2)$. Letting $cx \geq d$ and $Ax \geq b$ stand for the appropriate inequality systems in Pudlák's exponentiality result for Gomory-Chvátal cutting-plane proofs, the results in Dash [3] imply that $\sum_{i=1}^L t_i^2$ is exponential in the number of variables and constraints in $Ax \geq b$. Therefore, $\sum_{i=1}^L t_i$ – the total number of terms in mixing inequalities used in the proof – is exponential in the encoding size of $Ax \geq b$. \square

REFERENCES

- [1] V. Chvatal, Edmonds Polytopes and a hierarchy of combinatorial problems, *Discrete Mathematics* **4** 305–337 (1973).
- [2] W. J. Cook, R. Kannan, and A. Schrijver, Chvátal closures for mixed integer programming problems, *Mathematical Programming* **47** 155–174 (1990).
- [3] S. Dash. On the complexity of cutting plane proofs using split cuts. IBM Research Report RC24082, Oct. 2006.
- [4] S. Dash. Mixed integer rounding cuts and master group polyhedra. IBM Research Report RC24521, March 2008.
- [5] S. Dash, O. Günlük and A. Lodi. On the MIR closure of polyhedra, *Mathematical Programming* .to appear
- [6] S. S. Dey, A Note on Split Rank of Intersection Cuts, CORE DP 56, 2008.
- [7] O. Günlük, and Y. Pochet, Mixing Mixed-Integer Inequalities, *Mathematical Programming* **90** 429–457 (2001).
- [8] G. Nemhauser and L. A. Wolsey, *Integer and Combinatorial Optimization*, Wiley, New York (1988).
- [9] G. Nemhauser and L. A. Wolsey, A recursive procedure to generate all cuts for 0-1 mixed integer programs, *Mathematical Programming* **46**, 379–390 (1990).
- [10] P. Pudlák, Lower bounds for resolution and cutting plane proofs and monotone computations, *Journal of Symbolic Logic* **62** 981–998 (1997).
- [11] L.A. Wolsey, *Integer Programming*, Wiley, New York (1998).

Appendix. We now show how to derive $mix1_{\{1, \dots, n\}}$ as an MIR inequality (1.8) from the inequalities $s + z_1 \geq b_1$, $s + b_1 z_1 \geq b_1$, $mix1_{\{2, \dots, n\}}$ and $mix1_{\{1, k, \dots, n\}}$ for $k = 3, \dots, n$. Towards this end, we convert these inequalities to equations by adding

non-negative slack variables in the following manner:

$$s + z_1 - v_1 = b_1, \quad (M_1)$$

$$s + b_1 z_1 - v_2 = b_1, \quad (M_2)$$

$$s + b_1 z_1 + (b_k - b_1) z_k + \sum_{i=k+1}^n (b_i - b_{i-1}) z_i - v_k = b_n \quad (k = 3, \dots, n), \quad (M_k)$$

$$s + b_2 z_2 + \sum_{i=3}^n (b_i - b_{i-1}) z_i - v_{n+1} = b_n. \quad (M_{n+1})$$

Note that (M_1) - (M_{n+1}) involve $n - 1$ or fewer variables from z_1, \dots, z_n .

We define a multiplier λ_k for inequality (M_k) for $k = 1, \dots, n + 1$ and use these multipliers to obtain a base inequality (1.7) such that the MIR inequality (1.8) equals inequality (1.1). First, let

$$\mu_k = \begin{cases} (b_2 - b_1) \left(\frac{1}{1 - b_1} \right) & \text{if } k = 1, \\ (b_2 - b_1) \left(\frac{1}{b_n - b_1} - \frac{1}{1 - b_1} \right) & \text{if } k = 2, \\ (b_2 - b_1) \left(\frac{1}{b_{k-1} - b_1} - \frac{1}{b_k - b_1} \right) & \text{for } k = 3, \dots, n, \\ -1 & \text{if } k = n + 1, \end{cases}$$

and note that

$$\sum_{k=3}^p \mu_k = (b_2 - b_1) \sum_{k=3}^p \left(\frac{1}{b_{k-1} - b_1} - \frac{1}{b_k - b_1} \right) = 1 - \frac{b_2 - b_1}{b_p - b_1}.$$

Furthermore

$$\sum_{k=1}^n \mu_k = \frac{b_2 - b_1}{1 - b_1} + (b_2 - b_1) \left(\frac{1}{b_n - b_1} - \frac{1}{1 - b_1} \right) + 1 - \frac{b_2 - b_1}{b_n - b_1} = 1.$$

Now consider $\sum_{k=1}^n \mu_k (M_k)$ and denote it by

$$\alpha_0 s + \sum_{k=1}^n \alpha_k z_k - \sum_{k=1}^n \mu_k v_k = \beta. \quad (5.1)$$

Note that $\alpha_0 = \sum_{k=1}^n \mu_k = 1$, and $\alpha_2 = 0$. In addition,

$$\alpha_1 = \mu_1 + b_1 \sum_{k=2}^n \mu_k = \mu_1 (1 - b_1) + b_1 \sum_{k=1}^n \mu_k = (1 - b_1) \frac{b_2 - b_1}{1 - b_1} + b_1 = b_2.$$

For $k = 3, \dots, n$

$$\begin{aligned}
\alpha_k &= \sum_{l=3}^{k-1} \mu_l (b_k - b_{k-1}) + \mu_k (b_k - b_1) \\
&= (b_k - b_{k-1}) \left(1 - \frac{b_2 - b_1}{b_{k-1} - b_1}\right) + (b_2 - b_1) \left(\frac{1}{b_{k-1} - b_1} - \frac{1}{b_k - b_1}\right) (b_k - b_1) \\
&= (b_k - b_{k-1}) - \frac{(b_k - b_{k-1})(b_2 - b_1)}{b_{k-1} - b_1} + \frac{(b_2 - b_1)(b_k - b_1)}{b_{k-1} - b_1} - (b_2 - b_1) \\
&= b_k - b_{k-1}.
\end{aligned}$$

Finally,

$$\beta = b_n \sum_{k=3}^n \mu_k + b_1(\mu_1 + \mu_2) = b_n \left(1 - \frac{b_2 - b_1}{b_n - b_1}\right) + b_1 \frac{b_2 - b_1}{b_n - b_1} = b_n - (b_2 - b_1)$$

and therefore equation (5.1) is the same as

$$s + b_2 z_1 + \sum_{k=3}^n (b_k - b_{k-1}) z_k - \sum_{k=1}^n \mu_k v_k = b_n - (b_2 - b_1). \quad (5.2)$$

Further, $\sum_{i=1}^{n+1} \mu_k M_k$ equals

$$b_2(z_1 - z_2) + v_{n+1} - \sum_{k=1}^n \mu_k v_k = -(b_2 - b_1).$$

Therefore, if we define $\lambda_k = \mu_k/b_2$, and drop terms with negative coefficients for v_k variables in $\sum_{i=1}^{n+1} \lambda_k M_k$ we get

$$(z_1 - z_2) + v_{n+1}/b_2 \geq -(b_2 - b_1)/b_2.$$

If we let γ stand for the right-hand-side of the inequality above, then $\hat{\gamma} = b_1/b_2$. Applying the basic mixed-integer inequality, we get $(b_1/b_2)(z_1 - z_2) + v_{n+1}/b_2 \geq 0$ or

$$b_1(z_1 - z_2) + v_{n+1} \geq 0$$

as an MIR inequality. Substituting out v_{n+1} in the previous inequality using (M_{n+1}) , we get $mix1_{\{1, \dots, n\}}$.