

HIGH ACCURACY SEMIDEFINITE PROGRAMMING BOUNDS FOR KISSING NUMBERS

HANS D. MITTELMANN AND FRANK VALLENTIN

ABSTRACT. The kissing number in n -dimensional Euclidean space is the maximal number of non-overlapping unit spheres which simultaneously can touch a central unit sphere. Bachoc and Vallentin developed a method to find upper bounds for the kissing number based on semidefinite programming. This paper is a report on high accuracy calculations of these upper bounds for $n \leq 24$. The bound for $n = 16$ implies a conjecture of Conway and Sloane: There is no 16-dimensional periodic sphere packing with average theta series

$$1 + 7680q^3 + 4320q^4 + 276480q^5 + 61440q^6 + \dots$$

1. INTRODUCTION

In geometry, the *kissing number* in n -dimensional Euclidean space is the maximal number of non-overlapping unit spheres which simultaneously can touch a central unit sphere. The kissing number is only known in dimensions $n = 1, 2, 3, 4, 8, 24$, and there were many attempts to find good lower and upper bounds. We refer to Casselman [4] for the history of this problem and to Pfender, Ziegler [14], Elkies [7], and Conway, Sloane [6] for more background information on sphere packing problems.

Bachoc and Vallentin [1] develop a method (Section 2 recalls it) to find upper bounds for the kissing number based on semidefinite programming. Table 1 in Section 3, the main contribution of this paper, gives the values, i.e. the first ten significant digits, of these upper bounds for all dimensions $3, \dots, 24$. In all cases they are the best known upper bounds. Dimension 5 is the first dimension in which the kissing number is not known. With our computation we could limit the range of possible values from $\{40, \dots, 45\}$ to $\{40, \dots, 44\}$. In Section 4 we show that the high accuracy computations for the upper bounds in dimension 4 result into a question about a possible approach to prove the uniqueness of the kissing configuration in 4 dimensions.

Although acquiring the data for the table is a purely computational task we think that providing this table is valuable for several reasons: The kissing number is an important constant in geometry and results can depend on good upper bounds for it. For instance in Section 5 we show that there is no periodic point set in dimension

Date: May 4, 2009.

1991 Mathematics Subject Classification. 11F11, 52C17, 90C10.

Key words and phrases. kissing number, semidefinite programming, average theta series, extremal modular form.

The second author was partially supported by the Deutsche Forschungsgemeinschaft (DFG) under grant SCHU 1503/4.

16 with average theta series

$$1 + 7680q^3 + 4320q^4 + 276480q^5 + 61440q^6 + \dots$$

This proves a conjecture of Conway and Sloane [6, Chapter 7, page 190]. Furthermore, the actual computation of the table was very challenging. Bachoc and Vallentin [1] gave results for dimension $3, \dots, 10$. However, they report on numerical difficulties which prevented them from extending their results. Now using new, more sophisticated high accuracy software and faster computers and more computation time we could overcome some of the numerical difficulties. Section 3 contains details about the computations.

2. NOTATION

In this section we set up the notation which is needed for our computation. For more information we refer to [1]. For natural numbers d and $n \geq 3$ let $s_d(n)$ be the optimal value of the minimization problem

$$\begin{aligned} \min \left\{ 1 + \sum_{k=1}^d a_k + b_{11} + \langle F_0, S_0^n(1, 1, 1) \rangle : \right. \\ a_1, \dots, a_d \in \mathbb{R}, \quad a_1, \dots, a_d \geq 0, \\ b_{11}, b_{12}, b_{22} \in \mathbb{R}, \quad \begin{pmatrix} b_{11} & b_{12} \\ b_{12} & b_{22} \end{pmatrix} \text{ is positive semidefinite,} \\ F_k \in \mathbb{R}^{(d+1-k) \times (d+1-k)}, \quad F_k \text{ is positive semidefinite, } \quad k = 0, \dots, d, \\ q, q_1 \in \mathbb{R}[u], \quad \deg(p + pq_1) \leq d, \quad p, p_1 \text{ sums of squares,} \\ r, r_1, \dots, r_4 \in \mathbb{R}[u, v, t], \quad \deg\left(r + \sum_{i=1}^4 p_i r_i\right) \leq d, \quad r, r_1, \dots, r_4 \text{ sums of squares,} \\ 1 + \sum_{k=1}^d a_k P_k^n(u) + 2b_{12} + b_{22} + 3 \sum_{k=0}^d \langle F_k, S_k^n(u, u, 1) \rangle + q(u) + p(u)q_1(u) = 0, \\ \left. b_{22} + \sum_{k=0}^d \langle F_k, S_k^n(u, v, t) \rangle + r(u, v, t) + \sum_{i=1}^4 p_i(u, v, t)r_i(u, v, t) = 0 \right\}. \end{aligned}$$

Here P_k^n is the normalized Jacobi polynomial of degree k with $P_k^n(1) = 1$ and parameters $((n-3)/2, (n-3)/2)$. In general, *Jacobi polynomials* with parameters (α, β) are orthogonal polynomials for the measure $(1-u)^\alpha(1+u)^\beta du$ on the interval $[-1, 1]$. Before we can define the matrices S_k^n we first define the entry (i, j) with $i, j \geq 0$ of the (infinite) matrix Y_k^n containing polynomials in the variables u, v, w by

$$\begin{aligned} (Y_k^n)_{i,j}(u, v, t) &= u^i v^j \cdot \\ & \left((1-u^2)(1-v^2) \right)^{k/2} P_k^{n-1} \left(\frac{t-uv}{\sqrt{(1-u^2)(1-v^2)}} \right). \end{aligned}$$

Then we get S_k^n by symmetrization: $S_k^n = \frac{1}{6} \sum_{\sigma} \sigma Y_k^n$, where σ runs through all permutations of the variables u, v, t which acts on the matrix coefficients in the

obvious way. The polynomials p, p_1, \dots, p_4 are given by

$$\begin{aligned} p(u) &= -(u+1)(u+1/2), \\ p_1(u, v, t) &= p(u), \quad p_2(u, v, t) = p(v), \quad p_3(u, v, t) = p(t), \\ p_4(u, v, t) &= 1 + 2uvt - u^2 - v^2 - t^2. \end{aligned}$$

By $\langle A, B \rangle$ we denote the inner product between symmetric matrices $\text{trace}(AB)$.

In [1] it is shown that this minimization problem is a semidefinite program and that every upper bound on $s_d(n)$ provides an upper bound for the kissing number in dimension n . Clearly, the numbers $s_d(n)$ form a monotonic decreasing sequence in d .

3. BOUNDS FOR KISSING NUMBERS

n	best lower bound known	best upper bound previously known	SDP bound
3	12	<u>12</u> [16] Schütte, v.d. Waerden, 1953	$s_{11}(3) = 12.42167009 \dots$ $s_{12}(3) = 12.40203212 \dots$ $s_{13}(3) = 12.39266509 \dots$ $s_{14}(3) = 12.38180947 \dots$
4	24	<u>24</u> [11] Musin, 2008	$s_{11}(4) = 24.10550859 \dots$ $s_{12}(4) = 24.09098111 \dots$ $s_{13}(4) = 24.07519774 \dots$ $s_{14}(4) = 24.06628391 \dots$
5	40	45 [1] Bachoc, Vallentin, 2008	$s_{11}(5) = 45.06107293 \dots$ $s_{12}(5) = 45.02353644 \dots$ $s_{13}(5) = 45.00650838 \dots$ $s_{14}(5) = \underline{44.99899685} \dots$
6	72	<u>78</u> [1] Bachoc, Vallentin, 2008	$s_{11}(6) = 78.58344077 \dots$ $s_{12}(6) = 78.35518719 \dots$ $s_{13}(6) = 78.29404232 \dots$ $s_{14}(6) = 78.24061272 \dots$
7	126	135 [1] Bachoc, Vallentin, 2008	$s_{11}(7) = \underline{134.8824614} \dots$ $s_{12}(7) = 134.7319671 \dots$ $s_{13}(7) = 134.5730609 \dots$ $s_{14}(7) = 134.4488169 \dots$
8	240	<u>240</u> [12] Odlyzko, Sloane, 1979 [9] Levenshtein, 1979	$s_{11}(8) = 240.0000000 \dots$
9	306	366 [1] Bachoc, Vallentin, 2008	$s_{11}(9) = 365.3229274 \dots$ $s_{12}(9) = \underline{364.7282746} \dots$ $s_{13}(9) = 364.3980087 \dots$ $s_{14}(9) = 364.0919287 \dots$
10	500	567 [1] Bachoc, Vallentin, 2008	$s_{11}(10) = 558.1442813 \dots$ $s_{12}(10) = 556.2840736 \dots$ $s_{13}(10) = 555.2399024 \dots$ $s_{14}(10) = \underline{554.5075418} \dots$
11	582	915 [12] Odlyzko, Sloane, 1979	$s_{11}(11) = 878.6158044 \dots$ $s_{12}(11) = 873.3790094 \dots$ $s_{13}(11) = 871.9718533 \dots$ $s_{14}(11) = \underline{870.8831157} \dots$

12	840	1416 [12] Odlyzko, Sloane, 1979	$s_{11}(12) = 1364.683810 \dots$ $s_{12}(12) = 1362.200297 \dots$ $s_{13}(12) = 1359.283834 \dots$ $s_{14}(12) = \underline{1357.889300} \dots$
13	1130	2233 [12] Odlyzko, Sloane, 1979	$s_{11}(13) = 2089.116331 \dots$ $s_{12}(13) = 2080.631518 \dots$ $s_{13}(13) = 2073.074796 \dots$ $s_{14}(13) = \underline{2069.587585} \dots$
14	1582	3492 [12] Odlyzko, Sloane, 1979	$s_{11}(14) = 3224.950751 \dots$ $s_{12}(14) = 3202.448902 \dots$ $s_{13}(14) = 3189.127644 \dots$ $s_{14}(14) = \underline{3183.133169} \dots$
15	2564	5431 [12] Odlyzko, Sloane, 1979	$s_{11}(15) = 4949.650431 \dots$ $s_{12}(15) = 4893.479446 \dots$ $s_{13}(15) = 4876.037229 \dots$ $s_{14}(15) = \underline{4866.245659} \dots$
16	4320	8312 [13] Pfender, 2007	$s_{11}(16) = 7515.952644 \dots$ $s_{12}(16) = 7432.720718 \dots$ $s_{13}(16) = 7374.093742 \dots$ $s_{14}(16) = \underline{7355.809036} \dots$
17	5346	12210 [13] Pfender, 2007	$s_{11}(17) = 11568.41674 \dots$ $s_{12}(17) = 11333.84265 \dots$ $s_{13}(17) = 11128.26227 \dots$ $s_{14}(17) = \underline{11072.37543} \dots$
18	7398	17877 [12] Odlyzko, Sloane, 1979	$s_{11}(18) = 17473.48016 \dots$ $s_{12}(18) = 17034.32488 \dots$ $s_{13}(18) = 16686.28908 \dots$ $s_{14}(18) = \underline{16572.26478} \dots$
19	10668	25900 [3] Boyvalenkov, 1994	$s_{11}(19) = 26397.34794 \dots$ $s_{12}(19) = 25636.98958 \dots$ $s_{13}(19) = 25029.87432 \dots$ $s_{14}(19) = \underline{24812.30254} \dots$
20	17400	37974 [12] Odlyzko, Sloane, 1979	$s_{11}(20) = 39045.32761 \dots$ $s_{12}(20) = 37844.10380 \dots$ $s_{13}(20) = 37067.18966 \dots$ $s_{14}(20) = \underline{36764.40138} \dots$
21	27720	56851 [3] Boyvalenkov, 1994	$s_{11}(21) = 58087.03857 \dots$ $s_{12}(21) = 56079.21685 \dots$ $s_{13}(21) = 55170.03449 \dots$ $s_{14}(21) = \underline{54584.76757} \dots$
22	49896	86537 [12] Odlyzko, Sloane, 1979	$s_{11}(22) = 87209.06261 \dots$ $s_{12}(22) = 84922.09101 \dots$ $s_{13}(22) = 84117.92103 \dots$ $s_{14}(22) = \underline{82340.08003} \dots$
23	93150	128095 [3] Boyvalenkov, 1994	$s_{11}(23) = 128360.7969 \dots$ $s_{12}(23) = 127323.7095 \dots$ $s_{13}(23) = 125978.7655 \dots$ $s_{14}(23) = \underline{124416.9796} \dots$
24	196560	<u>196560</u> [12] Odlyzko, Sloane, 1979 [9] Levenshtein, 1979	$s_{11}(24) = 196560.0000 \dots$

TABLE 1. New upper bounds for the kissing number (best known values are underlined).

Finding the solution of the semidefinite program defined in Section 2 is a computational challenge. It turns out that the major obstacle is numerical instability and not the problem size. When d is fixed, then the size of the input matrices stays constant with n ; when n is fixed, then it grows rather moderately with d .

There is a number of available software packages for solving semidefinite programs. Mittelmann compares many existing packages in [10]. For our purpose first order, gradient-based methods such as SDPLR are far too inaccurate, and second order, primal-dual interior point methods are more suitable. Here increasingly ill-conditioned linear systems have to be solved even if the underlying problem is well-conditioned. This happens in the final phase of the algorithm when one approaches an optimal solution. Our problems are not well-conditioned and even the most robust solver `SeDuMi` which uses partial quadruple arithmetic in the final phase does not produce reliable results for $d > 10$.

We thus had to fall back on the implementation `SDPA-GMP` [8] which is much slower but much more accurate than other software packages because it uses the GNU Multiple Precision Arithmetic Library. We worked with 200–300 binary digits and relative stopping criteria of 10^{-30} . The ten significant digits listed in the table are thus guaranteed to be correct. One problem was the convergence. Even with the control parameter settings recommended by the authors of `SDPA-GMP` for “`slow but stable`” computations, the algorithm failed to converge in several instances. However, we found parameter choices which worked for all cases: We varied the parameter `lambdaStar` between 100 and 10000 depending on the case while the other parameters could be chosen at or near the values recommended for “`slow but stable`” performance.

The computations were done on Intel Core 2 platforms with one and two Quad processors. The computation time varied between five and ten weeks per case for $d = 12$. An accuracy of 128 bits in `SDPA-GMP` did yield sufficient accuracy but did not yield a reduction in computing time.

After the computations for the cases $d = 11$ and $d = 12$ were finished new 128-bit versions (quadruple precision) of `SDPA` and `CSDP` became available; partly with our assistance. These new versions do not rely on the GNU Multiple Precision Arithmetic Library. So the computation time for the cases $d = 13$ and $d = 14$ were reasonable: from one week to two and a half weeks.

4. QUESTION ABOUT THE OPTIMALITY OF THE D_4 ROOT SYSTEM

Looking at the values $s_d(4)$ in Table 1 one is led to the following question:

Question 4.1. *Is $\lim_{d \rightarrow \infty} s_d(4) = 24$?*

If the answer to this question is yes, then it would have two noteworthy consequences about optimality properties of the root system D_4 .

The root system D_4 defines (up to orthogonal transformations) a point configuration on the unit sphere $S^3 = \{x \in \mathbb{R}^4 : x \cdot x = 1\}$ consisting of 24 points; it is the same point configuration as the one coming from the vertices of the regular 24 cell. This point configuration has the property that the spherical distance of every two distinct points is at least $\arccos 1/2$. Hence, these points can be the maximal 24 touching points of unit spheres kissing the central unit sphere S^3 .

If $\lim_{d \rightarrow \infty} s_d(4) = 24$, then this would prove that the root system D_4 is the unique optimal point configuration of cardinality 24. Here optimality means that

one cannot distribute 24 points on S^3 so that the minimal spherical distance between two distinct points exceeds $\arccos 1/2$. Thus, the root system D_4 would be characterized by its kissing property. This is generally believed to be true but so far no proof could be given.

Another consequence would be that there is no universally optimal point configuration of 24 points in S^3 as conjectured by Cohn, Conway, Elkies, Kumar [5]. Universally optimal point configurations minimize every absolutely monotonic potential function. The conjecture would follow if the answer to our question is yes: Every universally optimal point configuration is automatically optimal and Cohn, Conway, Elkies, Kumar [5] show that the root system D_4 is not universally optimal.

5. NONEXISTENCE OF A SPHERE PACKING

Our new upper bound of 7355 for the kissing number in dimension 16 implies that there is no periodic point set in dimension 16 whose average theta series equals

$$(1) \quad 1 + 7680q^3 + 4320q^4 + 276480q^5 + 61440q^6 + \dots$$

This settles a conjecture of Conway and Sloane [6, Chapter 7, page 190]. In this section we explain this result. We refer to Conway, Sloane [6], Elkies [7], and to Bowert [2] for more information.

An n -dimensional *periodic point set* Λ is a finite union of translates of an n -dimensional lattice, i.e. one can write Λ as $\Lambda = (AZ^n + v_1) \cup \dots \cup (AZ^n + v_N)$, with $v_1, \dots, v_N \in \mathbb{R}^n$, and $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a linear isomorphism. The *average theta series* of Λ is

$$\Theta_\Lambda(z) = \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^N \sum_{v \in \mathbb{Z}^n} q^{\|Av - v_i + v_j\|^2}, \text{ with } q = e^{\pi iz}.$$

This is a holomorphic function defined on the complex upper half-plane. A holomorphic function f which is defined on the complex upper half-plane, which is meromorphic for $z \rightarrow i\infty$, and which satisfies the transformation laws

$$f(-1/z) = z^8 f(z), \text{ and } f(z+2) = f(z) \text{ for all } z \in \mathbb{C} \text{ with } \Im z > 0,$$

is called a *modular form* of weight 8 for the Hecke group $G(2)$. The expression (1) defines the unique modular form of weight 8 for the Hecke group $G(2)$ which starts off with $1 + 0q^1 + 0q^2$. It is also called an *extremal modular form*, see Scharlau and Schulze-Pillot [15]

If there would be a 16-dimensional periodic point set whose average theta series coincides with (1) then this periodic point set would define the sphere centers of a sphere packing with extraordinary high density (see [6, Chapter 7, page 190]). At the same time the existence of such a periodic point set would show that the kissing number in dimension 16 is at least 7680. This is not the case.

ACKNOWLEDGEMENTS

We thank Etienne de Klerk and Renata Sotirov for initiating our collaboration and we thank Frank Bowert and Rudolf Scharlau for bringing the conjecture of Conway and Sloane to our attention.

REFERENCES

- [1] C. Bachoc, F. Vallentin, *New upper bounds for kissing numbers from semidefinite programming*, J. Amer. Math. Soc. **21** (2008), 909–924.
- [2] F. Bowert, *Gewichtszähler und Distanzzähler von Codes und Kugelpackungen*, Ph.D. thesis, University of Dortmund, 2004.
- [3] P. Boyvalenkov, *Small improvements of the upper bounds of the kissing numbers in dimensions 19, 21 and 23*, Atti Sem. Mat. Fis. Univ. Modena **42** (1994), 159–163.
- [4] B. Casselman, *The difficulties of kissing in three dimensions*, Notices Amer. Math. Soc. **51** (2004), 884–885.
- [5] H. Cohn, J.H. Conway, N.D. Elkies, A. Kumar, *The D_4 root system is not universally optimal*, Exp. Math. **16** (2007), 313–320.
- [6] J.H. Conway, N.J.A. Sloane, *Sphere packings, lattices and groups*, third edition, Springer, 1999.
- [7] N.D. Elkies, *Lattices, linear codes, and invariants, Part I/II*, Notices Amer. Math. Soc. **47** (2000), 1238–1245, 1382–1391.
- [8] K. Fujisawa, M. Fukuda, K. Kobayashi, M. Kojima, K. Nakata, M. Nakata, M. Yamashita, *SDPA (SemiDefinite Programming Algorithm) and SDPA-GMP user’s manual — version 7.1.1*, Research Report B-448, Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, June 2008.
- [9] V.I. Levenshtein, *On bounds for packing in n -dimensional Euclidean space*, Soviet Math. Dokl. **20** (1979), 417–421.
- [10] H.D. Mittelmann, *An independent benchmarking of SDP and SOCP solvers*, Math. Prog. **95** (2003), 407–430.
- [11] O.R. Musin, *The kissing number in four dimensions*, Ann. of Math. **168** (2008), 1–32.
- [12] A.M. Odlyzko, N.J.A. Sloane, *New bounds on the number of unit spheres that can touch a unit sphere in n dimensions*, J. Combin. Theory Ser. A **26** (1979), 210–214.
- [13] F. Pfender, *Improved Delsarte bounds for spherical codes in small dimensions*, J. Combin. Theory Ser. A **114** (2007), 1133–1147.
- [14] F. Pfender, G.M. Ziegler, *Kissing numbers, sphere packings and some unexpected proofs*, Notices Amer. Math. Soc. **51** (2004), 873–883.
- [15] R. Scharlau, R. Schulze-Pillot, *Extremal lattices*, p. 139–170 in *Algorithmic algebra and number theory (Heidelberg, 1997)*, Springer, 1999.
- [16] K. Schütte, B.L. van der Waerden, *Das Problem der dreizehn Kugeln*, Math. Ann. **125** (1953) 325–334.

H.D. MITTELMANN, SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARIZONA STATE UNIVERSITY, TEMPE, AZ 85287-1804, USA
E-mail address: `mittelmann@asu.edu`

F. VALLENTIN, DELFT INSTITUTE OF APPLIED MATHEMATICS, TECHNICAL UNIVERSITY OF DELFT, P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS
E-mail address: `f.vallentin@tudelft.nl`